

FOR NON-ORAL HEALTH PROFESSIONALS

Dentine Hypersensitivity

Objectives

The aim of this fact sheet is to provide non-oral health professionals with an understanding of Dentine Hypersensitivity (DH) to include its:

- background
- aetiology and predisposing factors
- diagnosis
- management



Background

Dentine is the mineralized dental tissue layer that makes up the bulk of a tooth. It lies beneath the hard enamel of the tooth's crown and extends below the gum line, encasing the roots of the tooth. A bone-like vital tissue, it has the capacity to respond to physiologic and pathologic stimuli.

Dentine hypersensitivity (DH), also referred to as tooth sensitivity, occurs when dentine becomes exposed to the oral environment, either due to enamel being lost on the tooth crown, and/or the result of gum recession. DH is a frequently encountered oral condition affecting up to half of the population¹, causing pain and discomfort. In some cases, it can lead to impaired quality of life^{2,3}. DH is characterized by a sharp, arresting, transient pain arising from exposed dentine, usually at the cervical margin of the tooth, just above the gum line. It is most commonly a response to cold drinks and foods, cold windy weather or, less commonly, sweet/sour and tactile stimuli^{2,4,5}.

It is believed that DH is caused by increased fluid movement inside microscopic dentinal tubules in healthy dentine, stimulating pain from tooth pulpal nerve endings; this is known as the hydrodynamic theory⁶.

There is a wide range of reported DH prevalence as a result of differing data capture methodology: a systematic review⁷ capturing data from questionnaires both with and without clinically diagnosed dentine hypersensitivity indicates 11.5% prevalence. Large recent general dental practice epidemiological studies with clinically diagnosed DH shows higher prevalence of up to 50% in adults populations^{1,8,9}. DH peaks between 38-47 years of age¹, with more females presenting with this oral condition than males¹⁰.

Aetiology and predisposing factors

DH occurs (Figure 1) when tooth dentine is exposed on the tooth root due to gingival recession (gum shrinkage)¹¹. This is caused by multifactorial aetiological associations, including combinations of periodontal (gum) diseases and their treatment, thin gingival phenotype, the position of the tooth in the bone and lack of bone covering the tooth root, oral piercings, smoking and traumatic toothbrushing^{4,11}. The other main associated aetiological agent is erosive tooth wear¹², removing enamel from the tooth crown and/or dentine from the root, due to frequent acid insult (intrinsic and extrinsic acids), often in combination with synergistic, abrasive tooth wear (traumatic toothbrushing). The dentinal tubules must be patent from the oral environment to the pulp tissues, erosive challenges being predominantly responsible for patency of dentinal tubules.




Figure 1: Cause of dentine hypersensitivity: dentinal tubules open to the oral environment due to enamel loss or gingival recession¹¹.

If the aetiological factors are not addressed and treatment not implemented, DH can become a long-term oral health problem, with most people experiencing DH for over two years². Symptoms often impact negatively on daily activities such as speaking, eating, drinking, and toothbrushing³.

Determining a diagnosis

DH is difficult to diagnose (Figure 2) because only a very small amount of dentine needs to be exposed to cause pain. Oral health professionals will want to exclude other conditions with similar pain symptoms such as gum disease, tooth decay, a cracked tooth or teeth, inadequate restorations, vital tooth bleaching, atypical facial pain, bruxism and poorly developed enamel and other enamel and dentine defects for example molar incisor hypomineralization (MIH)⁴.

It is important to note that pain associated with DH always occurs after exposure to an external stimulus, is transient in nature and is not a continuous or spontaneous pain². DH frequently occurs after an acidic drink or food stuff, with 85% of people with erosive tooth wear experiencing DH¹⁰. Conversations about DH do not routinely take place in dental surgeries which can lead to fewer diagnoses¹³. Patients felt that oral healthcare professionals only have a limited time available and that DH was subjective and less important than other conditions, therefore not worthy of discussion with the dental team¹³. Understanding the barriers to DH conversations and how to facilitate them will, for most people, lead to successful treatment to alleviate DH pain².

Dentine Hypersensitivity: Clinical manifestations and symptoms

Tooth structure	Exposed Dentine (can be very small area) Usually acj-amelocemental junction, rarely 1 tooth Buccal or lingual aspect (2:1)
Trigger	Cold/windy stimuli are worst
Pain	Short, sharp, transient pain on stimulation Not spontaneous pain, ceases on stimulus removal
Duration	Episodic, seasonal, majority of cases > 2 yrs, often takes many years to resolve
Prevalence	Variable, increasing in young and old age groups
Pulp status	Thought to be no pulpal pathology, 2min refractory period Transient pulpal inflammation
Exacerbates pain	Erosive challenge 85% patients with erosive tooth wear > DH Main DH risk factor is timing of intake of acid (increased sensitivity straight after acidic challenges)




Figure 2: Clinical Manifestations and Symptoms.

Management by non-oral healthcare professionals

- If the pain is thought to be dental in origin, that is short, sharp and only for the duration of the stimulus, which is usually a cold stimulus, refer to an oral healthcare professional
- An efficacious desensitising toothpaste could be recommended in the first instance (proven agents include stannous and arginine followed by calcium sodium phosphosilicates, nanohydroxyapatite, oxalate, potassium formulations)

Management by oral healthcare professional

- Start a conversation about DH.
- Identify main aetiological and predisposing factors with risk analysis.
- Give prevention advice to raise awareness of DH, empower the patient to modify aetiological factors, giving reassurance that most DH cases are treatable.
- Recommend regular, long-term application of proven, anti-sensitivity, home-use fluoride toothpaste¹⁴⁻¹⁶.
- If application of proven anti-sensitivity home use treatment is not effective, in-office professional treatment is available from an oral healthcare professional, e.g. prophylaxis pastes, varnishes and precipitants, resin-based materials for restorations, periodontal (gum) plastic surgery to cover the exposed root dentine, laser therapy^{14,17,18}.
- Follow up on regular basis.

Oral healthcare delivery framework

Ask

- Patient's complaint, pain history and impact on everyday life, pain characteristics and factors that trigger the pain
- Medical history
- Dental history
- Dietary history, to check for frequent acidic food and beverage consumption
- Toothbrushing habits, to check for aggressive or incorrect brushing technique.

Look

- Patient's response to cold stimulus on teeth, ask patient to identify teeth with pain and look for tooth wear and gum shrinkage.

Decide

- If the patient should be referred to an oral healthcare professional to determine the diagnosis and most appropriate treatment.

Act

- Once DH is diagnosed by an oral healthcare professional, reiterate oral hygiene education, encourage associated factor prevention and self-care in particular tooth wear from frequent erosive challenge and gum recession from traumatic toothbrushing, removal of oral piercings, stop smoking.
- Empower the patient to act, giving reassurance that most DH is treatable.
- Recommend regular long-term application of proven anti-sensitivity home use fluoride toothpaste for alleviation of DH.
- Encourage follow up on regular basis with oral healthcare professional

Document

- Pain history, symptoms, stimulating and relieving factors
- Medical history
- Dental history
- Risk factors such as consumption of acidic foods and drinks, vomiting/reflux habit, oral hygiene habits and bruxism
- Advice given to patient
- Referral to an oral healthcare professional.

This factsheet is supported by:



References

1. West NX, Davies M, Sculean A, et al. Prevalence of dentine hypersensitivity, erosive tooth wear, gingival recession and periodontal health in seven European countries. *J Dent.* 2024;150. Available from: doi.org/10.1016/j.jdent.2024.105364.
2. Pollard A, Khan I, Davies M, Claydon N, West NX. Comparative efficacy of self-administered dentifrices for the management of dentine hypersensitivity – A systematic review and network meta-analysis. *J Dent.* 2022;130. Available from: doi.org/10.1016/j.jdent.2023.104433.
3. Porritt JM, Sufi F, Barlow A, Baker SR. The role of illness beliefs and coping in the adjustment to dentine hypersensitivity. *J Clin Periodontol.* 2014;41(1):60-69. Available from: doi.org/10.1111/jcpe.12177.
4. West NE, West NX. Therapeutic Strategies for Erosive Tooth Wear related to Dentine Hypersensitivity. In: Schlueter N, Ganss C, Lussi A, eds. *Erosive Tooth Wear: From Diagnosis to Therapy.* Monographs in Oral Science. 3. Karger; 2025; p. 245-248. Available from: doi.org/10.1159/000543560
5. Gillam DG. Current diagnosis of dentin hypersensitivity in the dental office: an overview. *Clin Oral Investig.* 2013; Suppl1:S21-9. Available from: doi.10.1007/s00784-012-0911-1
6. M. Brännström. Sensitivity of dentine. *Oral Surg Oral Med Oral Pathol.* 1966;21(4): 517-526. Available from: doi.org/10.1016/0030-4220(66)90411-7
7. Favaro Zeola L, Soares PV, Cunha-Cruz J. Prevalence of dentin hypersensitivity: systematic review and meta-analysis. *J Dent.* 2019;81:1–6. Available from: doi. 10.1016/j.jdent.2022.104157.
8. Awad MA, El Kassas D, Al Harthi L. Dentine hypersensitivity and dentine exposure in Arab patient populations. *J Oral Rehabil.* 2020;47(4):473-479. Available from: doi.org/10.1111/joor.12927.
9. Wang ZY, Rong WS, Zhong YS et al. Prevalence of dentinal hypersensitivity and related factors in Chinese urban adults. *Zhonghua Kou Qiang Yi Xue Za Zhi (Chinese Journal of Stomatology).* 2024; 59(9):927-934. 2024;59(9):927-934. Available from: doi.10.3760/cma.j.cn112144-20231229-00312
10. West NX, Sanz M, Lussi A, Bartlett D, Bouchard P, Bourgeois D. Prevalence of dentine hypersensitivity and study of associated factors: a European population-based cross-sectional study. *J Dent.* 2013. Available from: doi.org/10.1016/j.jdent.2013.07.017
11. Kim Jw, Park J-C. Dentin hypersensitivity and emerging concepts for treatments. *J Oral Biosci.* 2017; 59(4). Available from: doi.org/10.1016/j.job.2017.09.001.
12. O'Toole S, Bartlett D. The relationship between dentine hypersensitivity, dietary acid intake and erosive tooth wear. *J Dent.* 2017 Dec;67:84-87. Available from: doi. 10.1016/j.jdent.2017.10.002.
13. Asimakopoulou K, West NX, Davies M, Gupta A, Parkinson C, Scambler S. Why don't patients with Dentine Hypersensitivity discuss the condition with their dentist? *J Dent* 150 105362. Available from: doi.org/10.1016/j.jdent.2024.105362
14. West N, Seong J and Davies M. Management of dentine hypersensitivity: efficacy of professionally and self-administered agents. *J of Clinical Periodontology* 42 (Suppl 16) ppS256-302 10.1111/jcpe.12336.
15. Behzadi S, Mohammadi Y, Rezaei-Soufi L, Farmany A. Occlusion effects of bioactive glass and hydroxyapatite on dentinal tubules: a systematic review. *Clin Oral Investig.* 2022 Oct;26(10):6061-6078. Available from: doi: 10.1007/s00784-022-04639-y.
16. Lin P-Y, Cheng Y-W, Chu C-Y, Chien K-L, Lin C-P, Tu Y-K. In-office treatment for dentin hypersensitivity: a systematic review and network meta-analysis. *J Clin Periodontol* 2013,40 53-64. <https://doi.org/10.1111/jcpe.12011>

17. Shabbir S, Ahmed S, Zaidi, S.J.A. et al. Efficacy of seventh generation bonding agents as desensitizers in patients with dentin hypersensitivity: a randomized clinical trial. BMC Oral Health 2024 24(1): 562. Available from: doi.org/10.1186/s12903-024-04352-0
18. Behniafar B, Noori F, Chiniforush N, Raei A. The effect of lasers in occlusion of dentinal tubules and reducing dentinal hypersensitivity, a scoping review. BMC Oral Health. 2024;24(1). Available from: [doi.10.1186/s12903-024-05182-w](https://doi.org/10.1186/s12903-024-05182-w)

Other Resources

FDI World Dental Federation. Policy statement on tooth wear. (<https://www.fdiworlddental.org/sites/default/files/2023->) (accessed February 9, 2025)

FDI World Dental Federation. Dentine Hypersensitivity (<https://www.fdiworlddental.org/dentin-hypersensitivity#>) (Accessed February 9, 2025)

Disclaimer:

The provided fact sheet offers general information and may require adaptation to suit the unique scope of work and regulations governing non-dental healthcare professionals in each country.

The Educational Module for Other Healthcare Professionals Project is supported by **HALÉON**